

Sulphite 10 T

M368

0.1 - 12 mg/L SO₃

DTNB

Instrument specific information

The test can be performed on the following devices. In addition, the required cuvette and the absorption range of the photometer are indicated.

Instrument Type	Cuvette	λ	Measuring Range
SpectroDirect, XD 7000, XD 7500	□ 10 mm	405 nm	0.1 - 12 mg/L SO ₃

Material

Required material (partly optional):

Reagents	Packaging Unit	Part Number
Sulfite LR	Tablet / 100	518020BT

Application List

- · Waste Water Treatment
- Galvanization

Notes

Variations in the length of the vial can extend the measuring range:

- 10 mm vial: 0.1 mg/L 10 mg/L, solution: 0.01
- 20 mm vial: 0.05 mg/L 5 mg/L, solution: 0.01
- 50 mm vial: 0.02 mg/L 2 mg/L, solution: 0.001






Determination of Sulphite with Tablet

Select the method on the device.

For this method, a ZERO measurement does not have to be carried out every time on the following devices: XD 7000, XD 7500




Fill 10 mm vial with sample.




Place **sample vial** in the sample chamber. • Pay attention to the positioning.




Press the **ZERO** button.



Remove **vial** from the sample chamber.



Empty vial.



Dry the vial thoroughly.

For devices that require no ZERO measurement, start here.



Put **10 mL sample** in the sample vessel.




Add SULFITE LR tablet.




Crush tablet(s) by rotating slightly and dissolve.






Fill 10 mm vial with sample.



Place **sample vial** in the sample chamber. • Pay attention to the positioning.



Press the **TEST** (XD: **START**)button.



Wait for 5 minute(s) reaction time.

Once the reaction period is finished, the measurement takes place automatically.

The result in mg/L Sulphite appears on the display.



Analyses

The following table identifies the output values can be converted into other citation forms.

Unit	Cite form	Scale Factor
mg/l	SO ₃ ²⁻	1
mg/l	Na ₂ SO ₃	1.5743

Chemical Method

DTNB

Appendix

Calibration function for 3rd-party photometers

Conc. = a + b•Abs + c•Abs² + d•Abs³ + e•Abs⁴ + f•Abs⁵

	□ 10 mm	
а	-4.72981 • 10 ⁻¹	
b	6.87211 • 10 ⁺⁰	
С		
d		
е		
f		

Bibliography

R.E. Humphrey, M.H. Ward, W. Hinze, Spectrophotometric determination of sulphite with 4,4'-dithio-dipyridine and 5,5'-dithiobis(2-nitrobenzoic acid), Anal. Chem., 1970, 42 (7), pp 698–702